

Phytochemical study of *Swertia longifolia*

*¹Hajimehdipoor H., ²Dijoux-Franca M.G., ²Mariotte A.M., ³Amanzadeh Y.,
⁴Sadat-Ebrahimi S.E., ⁵Ghazi-Khansari M. and ⁶Mozaffarian V.

¹Department of Herbal Drugs, Food and Drug Control Laboratories and Food and Drug Laboratory Research Center, MOH & ME, Tehran, Iran. ²UMR 5063 CNRS-UJF, Département de Pharmacochimie Moléculaire (DPM), Laboratoire de Pharmacognosie, Faculté de Pharmacie de Grenoble, France. ³Department of Pharmacognosy, ⁴Department of Medicinal Chemistry, Faculty of Pharmacy, ⁵Department of Pharmacology, Faculty of Medicine, Tehran University of Medical Sciences., ⁶Research Institute of Forests and Rangelands, Tehran, Iran.

Received 26 Apr 2008; Revised 16 Sep 2008; Accepted 20 Sep 2008

ABSTRACT

Background: *Swertia* spp. (Gentianaceae) grow widely in eastern and southern Asian countries such as Japan, China and India and are used as traditional remedy for gastrointestinal complains because of their bitter principles. Several studies have been carried out on hypoglycemic, hepatoprotective, mono amino oxidase inhibitory and antidepressant effects of these plants and it has been shown that xanthenes and iridoids are responsible for their activities.

Purpose of the study: In order to gain better knowledge of endemic plants of Flora Iranica, *Swertia longifolia* Boiss. growing in the northern parts of Iran, was subjected to phytochemical studies.

Methods: Dried and milled aerial parts of the plant were extracted with petroleum ether and ethanol of which results of petroleum ether extract has been reported previously. For purification of ethanol extract, it was acidified with acetic acid and subsequently extracted with chloroform and then with *n*-butanol. The *n*-butanol extract was analyzed using different chromatographic methods and the structures of the isolated components were established by means of spectroscopic techniques.

Results: Four components including an iridoid glycoside (loganic acid), a secoiridoid glycoside (gentiopicroside), a secoiridoid dilactone (gentiolactone) and a nucleoside (uridine) were isolated from *n*-butanol extract of the plant.

Major conclusion: Similar to other species of *Swertia*, iridoid and secoiridoid glycosides could be considered as major constituents of *Swertia longifolia* Boiss.

Keywords: *Swertia longifolia* Boiss., Gentianaceae, gentiolactone, gentiopicroside, loganic acid, uridine.

INTRODUCTION

The *Swertia* genus (Gentianaceae) comprises 170 species that grow abundantly in east and south Asia. Seventy nine species have been distributed in China of which about 20 species have been used in Chinese traditional medicine for treatment of hepatic, choleric and inflammatory diseases. Several species are also used in India and Pakistan. The plants of *Swertia* genus are rich sources of xanthenes, flavonoids, iridoid and secoiridoid glycosides and terpenoids (1). Many investigations have been carried out on pharmacological effects of *Swertia* species and some of their components in the pure form have been isolated. It has been demonstrated that they have hypoglycemic (2), hepatoprotective (3,4),

anti-HIV (5,6), antitubercular (7), mono amino oxidase inhibitory (8,9) and central nervous system depressant (10) activities. In Iran, two *Swertia* species exist: *S. lactea* and *S. longifolia* and the second one is endemic plant of Flora Iranica which grows in wet areas of Alborz mountains (north of Iran) (11). Last studies on this plant has shown that the aerial parts of plant contain xanthenes and xanthone glycosides (12,13). Another investigation has demonstrated hepatoprotective effects of methanol and petroleum ether extracts and swerchirin with xanthone structure as major component of the plant (3). In this investigation, extraction, purification and structure elucidation of iridoid and secoiridoid glycosides have been determined.

MATERIALS AND METHODS

Plant materials

Swertia longifolia Boiss. (aerial parts) was collected in July 2003 from the North of Iran, Mazandaran province, Lavashm mountains, ca. 2900 m, Hajimehdipoor, No. 81007 (TARI) and identified by Dr. V. Mozaffarian, Botanist, from Research Institute of Forests and Rangelands (Tehran).

General experimental procedures

¹H-NMR and ¹³C-NMR spectra were acquired on a Bruker Avance 400 NMR spectrometer with CDCl₃ or CD₃OD as solvents as appropriate and TMS as an internal standard. Chemical shifts are expressed in δ-values (ppm). COSY (Correlated spectroscopy), HMQC (Heteronuclear multiple quantum coherence) and HMBC (Heteronuclear multiple band correlation) spectra were obtained using the usual pulse sequences. EIMS (Electron ionization mass spectrometry) spectra were recorded on a Finnigan TSQ-Mat 70 (70 eV) spectrometer and DCIMS (Desorption chemical ionization mass spectrometry) spectra were recorded on a R210C quadripolar spectrometer. Column chromatography was performed using silica gel (Kieselgel 60, 0.063-0.2 mm, ASTM, Germany) and Sephadex LH-20 (25-100 μm, Pharmacia Fine Chemicals, Sweden). MPLC (Medium pressure liquid chromatography) was performed on a C₁₈ column (50-60 μm, MN, Germany) by MPLC apparatus of D-1000 Berlin 37. VLC (Vacuum liquid chromatography) was carried out with a silica gel column (35-70 μm, SDS, Germany). Separation by SPE (Solid phase extraction) was carried out using silica gel (35-70 μm, SDS, Germany). TLC (Thin layer chromatography) employed Merck Kieselgel 60F₂₅₄ and RP-18 WF₂₅₄ S (Germany).

Extraction and isolation

Dried and milled aerial parts of plant (1 kg) were extracted sequentially in a Soxhlet apparatus with petroleum ether (60-80 °C, 12 hrs) and ethanol (24 hrs) and the results of the analyses of Petroleum ether extract was reported previously (12). The ethanol extract of the defatted plant, was evaporated under reduced pressure and the residue was treated with aqueous 4% acetic acid (500 ml) and the mixture was filtered. The clarified aqueous acid fraction was partitioned with chloroform and then with *n*-butanol.

Isolation of compounds from *n*-butanol extract

n-Butanol extract was evaporated under reduced pressure (12 g) and subjected to a Sephadex LH-20 column (2.5×55 cm) and eluted with methanol

(1 l). This process yielded 5 fractions, of which two fractions were selected for further purification (fraction A, 8.95 g and B, 300.70 mg) since they contained major components as was determined by thin layer chromatography (TLC).

Purification of fraction A

Vacuum liquid chromatography (VLC) was performed on a silica gel column (7×3 cm). The solvents were CH₂Cl₂:EtOAc:MeOH 60:40:0, 60:20:20, 50:30:20, 30:50:20, 10:70:20, 0:80:20, 0:60:40, 0:50:50, 0:0:100. The volume was 350 ml for each step. The fractions obtained from CH₂Cl₂:EtOAc:MeOH 60:40:0 (fraction A₁, 42.4 mg) and 60:20:20 (fraction A₂, 1.13 g) were chosen for further purifications because they contained major components as was determined by TLC.

For separation of components of the fraction A₁, a solid phase extraction (SPE) method was used. The sample was subjected to a silica gel column (1.5×4.5 cm) and eluted with CH₂Cl₂:EtOAc:MeOH (each 24 ml) 100:0:0, 98:2:0, 96:4:0, 94:6:0, 92:8:0, 90:10:0, 88:12:0, 86:14:0, 84:16:0, 80:20:0, 76:24:0, 72:28:0, 68:32:0, 64:36:0, 60:40:0, 55:45:0, 50:50:0, 40:50:10, 30:50:20, 20:50:30, 10:50:40, 0:50:50, 0:40:60, 0:20:80, 0:0:100. The fractions obtained from CH₂Cl₂:EtOAc:MeOH 98:2:0 contained a secoiridoid dilactone 1 (1.1mg) (14,15).

Gentiolactone 1: Amorphous, C₁₀H₁₂O₅, MW: 212, IR ν_{max} (Cm⁻¹): 3500, 1745, 1725, 1600; ¹H-NMR (CDCl₃, 400 MHz): δ 1.00 (3H, t, J = 7.4 Hz, H-10), 1.71 (1H, dq, J = 14.4, 7.2 Hz, H-8a), 1.85 (1H, dq, J = 14.4, 7.2 Hz, H-8b), 2.64 (1H, m, H-6a), 2.81 (1H, m, H-6b), 4.49 (2H, m, H-7), 5.01 (1H, ddd, J = 16.4, 3.6, 2.4 Hz, H-3a), 5.25 (1H, dt, J = 16.4, 1.8 Hz, H-3b); ¹³C-NMR (CDCl₃, 100 MHz): 7.7 (C-10), 22.5 (C-6), 30.8 (C-8), 66.5 (C-7), 66.7 (C-3), 72.1 (C-9), 118.8 (C-4), 153.4 (C-5), 161.5 (C-11), 172.6 (C-1).

In order to separate components of the fraction A₂, this fraction was chromatographed over a silica gel column (3×39 cm) with CH₂Cl₂:EtOAc:MeOH 60:20:20, 50:20:30, 10:40:50, 0:0:100 as solvents. The volume was 830 ml for each step. Finally secoiridoid glycoside 2 (64.2 mg) (16) was obtained from CH₂Cl₂:EtOAc:MeOH 60:20:20 fractions and an iridoid glycoside 3 (57.2 mg) (17) was purified from CH₂Cl₂:EtOAc:MeOH 10:40:50 fractions.

Gentiopicroside 2: Amorphous, C₁₆H₂₀O₉, MW: 356, EIMS: *m/z* 357 (28), 195 (100), 176 (72); ¹H-NMR (CD₃OD, 400 MHz): δ 7.44 (1H, s, H-3), 5.75 (1H, ddd, J = 17.2, 10.4, 6.8 Hz, H-8), 5.66 (1H, d, J = 2.8 Hz, H-1), 5.61 (1H, m, H-6), 5.23




Figure 1. Structures of compounds 1-4

(1H, dt, $J = 15.6, 1.4$ Hz, H-10a), 5.19 (1H, dt, $J = 8.0, 1.4$ Hz, H-10b), 5.07 (1H, m, H-7a), 4.98 (1H, m, H-7b), 4.64 (1H, d, $J = 8.0$ Hz, H-1'), 3.89 (1H, dd, $J = 12.0, 2.4$ Hz, H-6'a), 3.64 (1H, dd, $J = 12.0, 6.4$ Hz, H-6'b), 3.35 (1H, t, $J = 8.8$ Hz, H-3'), 3.30 (2H, m, H-5', H-9), 3.23 (1H, t, $J = 8.8$ Hz, H-4'), 3.14 (1H, dd, $J = 9.2, 8.0$ Hz, H-2'); ^{13}C -NMR (CD_3OD , 100 MHz): δ 165.0 (C-11), 149.3 (C-3), 133.6 (C-8), 125.6 (C-5), 117.2 (C-10), 115.8 (C-6), 103.5 (C-4), 98.8 (C-1'), 97.1 (C-1), 77.0 (C-3'), 76.5 (C-5'), 73.1 (C-2'), 70.1 (C-4'), 69.6 (C-7), 61.3 (C-6'), 45.2 (C-9).

Loganic acid 3: Amorphous, $\text{C}_{16}\text{H}_{24}\text{O}_{10}$, MW: 376, DCIMS: m/z 394 (10), 232 (10), 215 (70), 212 (100), 198 (58), 180 (69), 153 (77), 145 (67), 127 (66); ^1H -NMR (CD_3OD , 400 MHz), δ 7.30 (1H, s, H-3), 5.20 (1H, d, $J = 4.4$ Hz, H-1), 4.60 (1H, d, $J = 8.0$ Hz, H-1'), 3.98 (1H, m, H-7), 3.83 (1H, dd, $J = 12.0, 1.6$ Hz, H-6'a), 3.61 (1H, dd, $J = 12.0, 5.6$ Hz, H-6'b), 3.32 (2H, m, H-5', H-3'), 3.24 (1H, m, H-4'), 3.15 (1H, dd, $J = 8.8, 8.0$ Hz, H-2'), 3.03 (1H, bq, $J = 8.0$ Hz, H-5), 2.17 (1H, ddd, $J = 14.0, 8.0, 1.2$ Hz, H-6 β), 1.95 (1H, dt, $J = 13.2, 4.0$ Hz, H-9), 1.80 (1H, m, H-8), 1.60 (1H, ddd, $J = 12.8, 8.0, 4.0$ Hz, H-6 α), 1.03 (1H, d, $J = 6.8$ Hz, H-10); ^{13}C -NMR (CD_3OD , 100 MHz), δ 170.1 (C-11), 150.3 (C-3), 113.3 (C-4), 98.6 (C-1'), 96.2 (C-1), 76.9 (C-5'), 76.6 (C-3'), 73.8 (C-

7), 73.3 (C-2'), 70.2 (C-4'), 61.3 (C-6'a), 45.1 (C-9), 41.3 (C-8), 40.7 (C-6), 30.8 (C-5), 12.1 (C-10).

Isolation of components from fraction B

MPLC was performed on a C_{18} column (1 \times 47 cm). The solvents were H_2O : MeOH 100:0 (220 ml), 95:5, 90:10, 85:15, 80:20, 70:30, 60:40, 50:50 (each 110 ml), 40:60 (160 ml), 30:70, 20:80, 10:90 (each 110 ml) and 0:100 (330 ml) using λ_{max} of 250 nm. The first fractions obtained from H_2O :MeOH 100:0 contained a pure nucleoside 4 (3.9 mg) (18,19).

Uridine 4: Amorphous, $\text{C}_9\text{H}_{12}\text{N}_2\text{O}_6$, MW: 244, EIMS: m/z 244 (4), 226 (23), 214 (27), 196 (39), 179 (100), 168 (78), 133 (54), 125 (88), 113 (57), 73 (66); ^1H -NMR (CD_3OD , 400 MHz): δ 7.95 (1H, d, $J = 8.0$ Hz, H-6), 5.84 (1H, d, $J = 4.4$ Hz, H-1'), 5.63 (1H, d, $J = 8.0$ Hz, H-5), 4.11 (1H, t, $J = 4.8$ Hz, H-2'), 4.08 (1H, t, $J = 4.8$ Hz, H-3'), 3.94 (1H, m, H-4'), 3.78 (1H, dd, $J = 12.0, 2.8$ Hz, H-5'a), 3.67 (1H, dd, $J = 12.0, 2.8$ Hz, H-5'b); ^{13}C -NMR (CD_3OD , 100 MHz): δ 164.8 (C-2), 151.1 (C-4), 141.3 (C-6), 101.2 (C-5), 89.3 (C-1'), 85.0 (C-4'), 74.3 (C-2'), 70.0 (C-3'), 60.9 (C-5').

RESULTS AND DISCUSSION

The family of Gentianaceae is apparently characterized by the universal occurrence of

iridoid glycosides. The iridoids which are found consistently are secoiridoids which occasionally result from carboxylic iridoids by biosynthetic pathway (20).

Preliminary examination of the *n*-butanol extract of the aerial parts of *Swertia longifolia* by analytical TLC suggested the presence of major components mainly with iridoid structure; which were isolated by chromatographic methods. Four components were purified from the extract and their structures determined by spectroscopic means. The spectroscopic data of compounds **1-4** were in good agreement with those described in the literature for gentiolactone (14,15), gentiopicroside (16), loganic acid (17) and uridine (18,19).

Uridine is a nucleoside with pyrimidine structure which is presents in all organisms as one of the RNA bases (21). Gentiopicroside is one of the most common secoiridoid glycosides presents in Gentianaceae which has been reported from 107 of 127 species in Gentianaceae, mainly in *Gentiana* nad *Swertia* genus (20). The compound has interesting pharmacological properties such as

anti-inflammatory (22), protective effects against many kinds of chemical and immunological liver injuries (23-25) and stress-induced gastric ulcers (26), bile secretion promoting property (25) and smooth muscle relaxing activity (27). It also exhibits cytoprotective effect which may cause a synergism in terms of wound healing activity of *Gentiana* root (28). Since gentiopicroside exists in *Swertia longifolia* as well, above-mentioned pharmacological properties could be expected from this plant. Loganic acid which is another compound of *S. longifolia*, has iridoid glycoside structure. It has been reported mainly from *Gentiana* species (29). However among *Swertia* species, loganic acid has been isolated only from *S. caroliniensis* (30, 31) and no report is available about occurrence of gentiolactone in *Swertia* species. Therefore, it is the first report of presence of this compound in *Swertia* genus. Although, several investigations have been performed on biological and pharmacological abilities of pure components isolated from plants such as iridoids and secoiridoids, no study has been carried out on effects of loganic acid and gentiolactone.

REFERENCES

1. Pant N, Jain DC, Bhakuni RS. Phytochemicals from genus *Swertia* and their biological activities. *Indian J chem* 2000; 39B: 565-586.
2. Basnet P, Kodota S, Shimizu M, Namba T. Bellidifolin: a potent hypoglycemic agent in streptozotocin-induced diabetic rats from *Swertia japonica*. *Planta Med* 1994; 60: 507-511.
3. Hajimehdipour H, Sadeghi Zh, Elmi S, Elmi A, Ghazi-Khansari M, Amanzadeh Y, Sadat-Ebrahimi SE. Protective effects of *Swertia longifolia* Boiss. and its active compound, swerchirin, on paracetamol-induced hepatotoxicity in mice. *J pharm Pharmacol* 2006; 58: 1-4.
4. Hase K, Li J, Basnet P, Xiong Q, Takamura Sh, Namba T, Kadota Sh. Hepatoprotective principles of *Swertia japonica* Makino on d-galactosamine/lipopolysaccharide-induced liver injury in mice. *Chem Pharm Bull* 1997; 45(11): 1823-1827.
5. Wang JN, Hou CY, Liu YL, Lin LZ, Gil RR, Cordell GA. Swertiflanchesides an HIV-reverse transcriptase inhibitor and the first flavone-xanthone dimer from *Swertia franchetiana*. *J Nat Prod* 1994; 57(2): 211-217.
6. Pengsuparp T, Cai L, Constant H, Fong HHS, Lin LZ, Kinghorn AD, Pezzuto JM, Cordell GA, Ingolfsson K, Wagner H, Hughes SH. Mechanistic evaluation of new plant-derived compounds that inhibit HIV-1 reverse transcriptase. *J Nat Prod* 1995; 58(7): 1024-1031.
7. Ghosal S, Biswas K, Chaudhuri RK. Chemical constituents of Gentianaceae XXIV: Antimycobacterium tuberculosis activity of naturally occurring xanthenes and synthetic analogs. *J Pharm Sci* 1978; 67(5): 721-722.
8. Hostettmann K, Wagner H. Xanthone glycosides. *Phytochemistry* 1977; 16: 821-829.
9. Bhattacharya SK, Ghosal S, Chaudhuri RK, Sanyal AK. *Conscora decussata* (Gentianaceae) xanthenes. III. Pharmacological studies. *J Pharm Sci* 1972; 61: 1838-1840.
10. Bhattacharya SK, Ghosal S, Chaudhuri RK, Singh AK, Sharma PV. Chemical constituents of Gentianaceae, XI, antipsychotic activity of gentianine. *J Pharm Sci* 1974; 63: 1341-1342.
11. Mozaffarian V. A dictionary of Iranian plant names. Tehran: Farhang Mo'aser; 1996, p. 530.
12. Hajimehdipour H, Amanzadeh Y, Sadat Ebrahimi SE, Mozaffarian V. Three tetraoxygenated xanthenes from *Swertia longifolia*. *Pharm Biol* 2003; 41: 497-499.
13. Hajimehdipour H, Dijoux-Franca MG, Mariotte AM, Amanzadeh Y, Sadat-Ebrahimi SE, Ghazi-Khansari M. Two new xanthone diglycosides from *Swertia longifolia* Boiss. *Nat Prod Res* 2006; 20: 1251-1257.
14. Suhr IH, Arends P, Jensen B. Gentiolactone, a secoiridoid dilactone from *Gentiana purpurea*. *Phytochemistry* 1978; 17: 135-138.

15. Tan RX, Kong LD. Secoiridoids from *Gentiana siphonantha*. *Phytochemistry* 1997; 46: 1035-1038.
16. Boros CA, Stermitz FR. Iridoids. An updated review. II. *J Nat Prod* 1991; 54: 1173-1246.
17. Boros CA, Stermitz FR. Iridoids. An updated review. I. *J Nat Prod* 1990; 53: 1055-1147.
18. Fromageot HPM, Griffin BE, Reese CB, Sulston JE, Trentham DR. Orientation of ribonucleoside derivatives by proton magnetic resonance spectroscopy. *Tetrahedron* 1966; 22: 705-710.
19. Kalinowski HO, Berger S, Braun S. Carbon-13 NMR spectroscopy. Chichester: John Wiley & Sons; 1991, p. 440.
20. Struwe L, Albert V. *Gentianaceae-systematics and natural history*. London: Cambridge University Press; 2002, p. 610.
21. Dey PM. *Plant Biochemistry*. London: Academic Press; 1997, pp. 483-484.
22. Chang-Xun Ch, Zhan-Wen L, Zheng-Rong S, Chun-Qing S, Zhi-Bi H. Studies on anti-inflammatory effect of gentiopicroside. *Zhongcaoyào* 2003; 34(9): 814-816.
23. Li YQ, Zhao DH, Pan BR, Li BG, Sun WJ, Tian Q, Jia M. Effects of gentiopicroside on liver injury of rats. *Journal of Fourth Military Medical University* 2001; 22: 1645-1649.
24. Kondo Y, Takano F, Hojo H. Suppression of chemically and immunologically induced hepatic injuries by gentiopicroside in mice. *Planta Med* 1994; 60: 414-416.
25. Liu ZW, Chen CX, Jin RM. Studies on liver-protecting and bilesecretion-promoting effects of gentiopicroside. *China Traditional and Herbal Drugs* 2002; 33: 47-50.
26. Niiho Y, Yamazaki T, Nakajima Y, Yamamoto T, Ando H, Hirai Y, Toriizuka K, Ida Y. Gastroprotective effects of bitter principles isolated from Gentian root and Swertia herb on experimentally-induced gastric lesions in rats. *J Nat Med* 2006; 60: 82-88.
27. Rojas A, Bah M, Rojas JI, Gutierrez DM. Smooth muscle relaxing activity of gentiopicroside isolated from *Gentiana spathacea*. *Planta Med* 2000; 66: 765-767.
28. Oztürk N, Korkmaz S, Oztürk Y, Başer KH. Effects of gentiopicroside, sweroside and swertiamarine, secoiridoids from gentian (*Gentiana lutea* ssp. *Symphyantra*), on cultured chicken embryonic fibroblasts. *Planta Med* 2006; 72(4): 289-294.
29. Ando H, Hirai Y, Fujii M, Hori Y, Fukumura M, Niiho Y, Nakajima Y, Shibata T, Toriizuka K, Ida Y. The chemical constituents of fresh Gentian root. *J Nat Med* 2007; 61(3): 269-279.
30. Coscia CJ, Guarnaccia R, Botta L. Monoterpene biosynthesis. I. Occurrence and mevalonoid origin of gentiopicroside and loganic acid in *Swertia carolinensis*. *Biochemistry* 1969; 8(12): 5036-5043.
31. Hill RA. *Dictionary of natural products*. London: Chapman & Hall; 1994, p. 3668.